



## Medicines That May Need To Be Adjusted (Talk With Your Physician)

May require less dose with smoking cessation			Possible Reason
Drug Name	Brand Name	Use/Type	
Acetaminophen	Tylenol Also active ingredient in: Percocet, Lortab, Lorcet, Ultracet, Esgic, Fioricet	Pain reliever; Also active ingredient in many combination prescription pain relievers	Deinduction of hepatic enzymes on smoking cessation
Caffeine	Excedrin, BC Powders, Esgic, Fioricet	Pain reliever, stimulant	
Imipramine	Tofranil	Antidepressant	
Pentazocine	Talwin	Pain reliever	
Propranolol	Inderal, Innopran XL	Blood Pressure/heart medication	
Theophylline	Theo-Dur, Slo-Bic, Uniphyll	Common group of breathing medications	
Insulin	Humulin, Novalin, Humalog, Novolog, Lantus	Commonly used in diabetics	
Adrenergic Antagonists eg., prazosin, labetalol	Mini-press, Normodyne Trandate, Hytrin	Used to reduce blood pressure, and treat BPH (prostate)	Increase in subcutaneous insulin absorption with smoking cessation
May require greater dose with smoking cessation			Possible Reasons
Drug Name	Brand Name	Use/Type	
Adrenergic Agonists eg., isoproterenol phenylephrine	Vospire, Proventil, Albuterol	Another group of commonly-used breathing medications	Decrease in circulating catecholamines with smoking cessation